

A Quick Look at Using DTaP/IPV/Hib (Pentacel®)

Indications for Use and Schedule

Approved for:

- Routine schedule of 2, 4, 6, and 15-18 months of age
- Ages 6 weeks through 4 years (4 years, 364 days)
- For doses 1, 2, 3 or 4 of DTaP, IPV or Hib

Do not use:

- For any person 5 years of age and older
- For the 5th dose of DTaP or, if needed, a 5th dose of IPV

Make sure minimum age and minimum intervals are met:

- Minimum age for dose 1 is 6 weeks
- Minimum age for dose 4 is 12 months
- Minimum intervals:
 - 4 weeks between dose 1 & 2
 - 4 weeks between dose 2 & 3
 - 6 months between dose 3 & 4


Vaccine Administration

- Intramuscular (IM) injection in the deltoid of the arm or anterolateral thigh
- 1 inch needle; 22-25 gauge
- Professional judgment is appropriate when selecting needle length and site
- Can be given with other vaccines, at the same visit

· See “Further Points” for reconstitution instructions

Storage and Handling

- Store the DTaP-IPV vial and the Hib vial in the original box and in the refrigerator at **35°-46° F (2°-8°C)**
- Shake well before using
- Administer within 30 min of reconstituting



CONTRAINDICATIONS

- An anaphylactic reaction to a prior dose of Pentacel®, DTaP, IPV or Hib vaccine
- An anaphylactic reaction to a component of DTaP/IPV/Hib (Pentacel) including neomycin or polymyxin B
- Encephalopathy not due to another cause occurring within 7 days after vaccination with a pertussis-containing vaccine
- Progressive neurologic disorder (such as infantile spasms, uncontrolled epilepsy or progressive encephalopathy)

PRECAUTIONS

- Moderate to severe acute illness
- Guillain-Barré syndrome within 6 weeks of a prior tetanus toxoid-containing vaccine
- Temperature greater than or equal to 105°F within 48 hours of vaccination with no other identifiable cause
- Collapse or shock-like state within 48 hours of vaccination
- Persistent inconsolable crying lasting more than 3 hours within 48 hours of vaccination
- Convulsions with or without a fever within 3 days of vaccination

FURTHER POINTS

- **Be sure to reconstitute the Hib vial with DTaP-IPV vial before administration. Use only this DTaP-IPV as the diluent.**
- CDC recommends that the lot numbers from the DTaP-IPV and the Hib vials be recorded on the vaccine administration record. Currently, MCIR will only allow for one lot number to be entered; use the lot number from the Hib vial (matches outer box #).
- Pentacel® does not contain a preservative. After removal of the dose, any remaining vaccine should be discarded.
- The combined DTaP-IPV-Hib vaccine may be used when any component of the vaccine is indicated, and if the other components are not contraindicated
- There is not a separate Vaccine Information Statement (VIS) for Pentacel. Use the current VISs for DTaP, IPV, and Hib or the Multi-vaccine VIS that include information about the Michigan Care Improvement Registry (MCIR).
- VISs with MCIR information are available at www.michigan.gov/immunize or your local health department
- Document as “DTaP/IPV/Hib” in MCIR, on the vaccine administration record & immunization record card

Publicly purchased DTaP/IPV/Hib (Pentacel) can be administered to eligible children 6 weeks through 4 years of age, utilizing the Vaccines for Children (VFC) Program, in private providers' offices. Eligible children include those who are uninsured, underinsured, Medicaid eligible, Native American or Alaskan Natives. Contact your local health department for more information

For additional information, refer to VFC Program Resolutions on “Vaccines to Prevent Diphtheria, Tetanus and Pertussis”, “Vaccines to Prevent Poliomyelitis”, and “Vaccines to Prevent *Haemophilus influenzae* type B” and the ACIP Recommendations on the use of DTaP, IPV and Hib vaccines, located at <http://www.cdc.gov/vaccines/recs>